arXiv:1105.1054v1 [math.GR] 5 May 2011

ON MAXIMAL SUBGROUPS OF A FINITE SOLVABLE GROUP

D. V. Gritsuk and V. S. Monakhov

January 20, 2013

Abstract

The following result is received: Let H be a non-normal maximal subgroup of a finite solvable group G and let $q \in \pi(F(H/\mathrm{Core}_G H))$, then G has a Sylow q-subgroup Q such that $N_G(Q) \subseteq H$.

Keywords: finite solvable group, Sylow subgroup, maximal subgroup

MSC2010 20D20, 20E34

1 Introduction

All groups considered in this paper will be finite. All notations and definitions correspond [1].

In 1986, V. A. Vedernikov obtained the following result.

Theorem A. [2, Corollary 2.1] *If H is a non-normal maximal subgroup of a solvable group G then $N_G(Q) \subseteq H$ for some Sylow subgroup Q of G.*

Here $N_G(Q)$ is the normalizer of Q in G.

In this note we consider the following question:

What is Sylow subgroup that its normalizer contained in non-normal maximal subgroup of solvable group?

The answer to this question was obtained in the following theorem.

Theorem 1. *Let H be a non-normal maximal subgroup of a solvable group G and let $q \in \pi(F(H/\mathrm{Core}_G H))$. Then G has a Sylow q-subgroup Q such that $N_G(Q) \subseteq H$.*

Here $F(X)$ is the Fitting subgroup of X, $\pi(Y)$ is the set of all prime divisors of $|Y|$, $\mathrm{Core}_G H = \cap_{g\in G} H^g$ is the core of H in G, i. e. the largest normal subgroup of G contained in H.

Corollary 1.1. *Let H be a non-normal maximal subgroup of a solvable group G and let $q \in \pi(F(H/\mathrm{Core}_G H))$. Then H has a Sylow q-subgroup Q such that $N_G(H_1) \subseteq H$ for each subgroup H_1 of H satisfying $Q \subseteq H_1 \subseteq H$.*

Corollary 1.2. *Let H be a non-normal maximal subgroup of a solvable group G and let $\omega \subset \pi(F(H/\mathrm{Core}_G H))$. Then G has a Hall ω-subgroup G_ω such that $N_G(G_\omega) \subseteq H$.*

For non-solvable groups, this result is false. For example, $PSL(2,17)$ has the order $2^4\cdot 3^2\cdot 17$ and the symmetric group S_4 is a maximal subgroup in $PSL(2,17)$, see [2]. Since $|S_4| = 2^3\cdot 3$, it follows that S_4 does not contain a Sylow subgroup of $PSL(2,17)$. Thus it is not possible to extend theorem of V. A. Vedernikov and Theorem 1 to the non-solvable groups.

The following question is contained in [2].

Question. (V. A. Vedernikov, 1986, [2]) *Can Theorem A be extended to a p-solvable group G containing a maximal subgroup M such that $|G : M| = p^a$?*

The positive answer to this question was obtained for $F(M/\mathrm{Core}_G M) \neq 1$ in the following theorem.

Theorem 2. *Let G be a p-solvable group. Let M be a non-normal maximal subgroup of G, and let $|G : M| = p^\alpha$. Then:*

1) *if $F(M/\mathrm{Core}_G M) \neq 1$ and $q \in \pi(F(M/\mathrm{Core}_G M))$, then G has a Sylow q-subgroup Q such that $N_G(Q) \subseteq M$;*

2) *if $F(M/\mathrm{Core}_G M) = 1$, then $N_G(K) \subseteq M$ for some Hall p'-subgroup K of G.*

2 Notations and preliminary results

In this section we give some definitions and basic results that will be used later in our paper.

Let $\mathbb{P}$ be the set of all prime numbers, and let π be the set of primes, i. e $\pi \subseteq \mathbb{P}$. In that paper, π' is the set of all primes not contained in π, i. e. $\pi = \mathbb{P} \setminus \pi'$, $\pi(m)$ is the set of prime divisors of m. If $\pi(m) \subseteq \pi$ then m is called *π-number*.

The subgroup H of G is called *π-subgroup* if $|H|$ is π-number. The subgroup H of G is called Hall *π-subgroup* if $|H|$ is π-number and $|G : H|$ is π'-number. As usually, $O_\pi(X)$ is the largest normal π-subgroup of X.

The group is called *π-separable* if it has a series whose factors are π-groups or π'-groups.

Lemma 1. [2, Theorem 1] *Let G be a π-separable group, and let H be a subgroup of G. If $|G : H|$ is π-number, then $O_\pi(H) \subseteq O_\pi(G)$.*

Lemma 2. *Let R be a Hall π-subgroup of a π-separable group G, and let N be a normal subgroup of G. Then $N_G(R)N/N = N_{G/N}(RN/N)$.*

Proof. For $x \in N_G(R)$ we obtain:

$$(x^{-1}N)(RN/N)(xN) = R^x N/N = RN/N,$$

i. e. $N_G(R)N/N \leq N_{G/N}(RN/N)$. Conversely, if $yN \in N_{G/N}(RN/N)$, then $R^y N = RN$. Now R and R^y are Hall subgroups of RN which are conjugate, i. e. $R^y = R^{ak} = R^k$ for some $ak \in RN$, $a \in R$, $k \in N$. Then $yk^{-1} \in N_G(R)$, whence $y \in N_G(R)N$, i. e. $N_{G/N}(RN/N) \leq N_G(R)N/N$. The Lemma is proved.

Lemma 3. *Let G be a π-solvable group containing a nilpotent Hall π-subgroup. If H is a maximal subgroup of G and $|G : H|$ is π-number, then $O_\pi(H)$ is a normal subgroup of G.*

Proof. By Lemma 1, $O_\pi(H) \subseteq O_\pi(G)$. If $O_\pi(H) = O_\pi(G)$ then $O_\pi(H)$ is a normal subgroup of G. Let $O_\pi(H)$ be a proper subgroup of $O_\pi(G)$, and let G_π be a Hall π-subgroup of G. Clearly, $O_\pi(G)$ is a proper subgroup of G_π. Since G_π is a nilpotent subgroup, we have $O_\pi(H)$ is a proper subgroup of $D = N_{G_\pi}(O_\pi(H))$. Since $O_\pi(H) = H \cap O_\pi(G)$, we have D does not contained in H, so $N_G(O_\pi(H)) \supseteq \langle H, D\rangle = G$ and $O_\pi(H)$ is a normal subgroup of G. The Lemma is proved.

3 Main results

Theorem 3. *Let G be a π-solvable group containing a nilpotent Hall π-subgroup. Let M be a non-normal maximal subgroup of G, and let $|G : M|$ is π-number. Then:*

1) if $F(M/\mathrm{Core}_G M) \neq 1$ and $q \in \pi(F(M/\mathrm{Core}_G M))$, then G has a Sylow q-subgroup Q such that $N_G(Q) \subseteq M$;

2) if $F(M/\mathrm{Core}_G M) = 1$, then $N_G(K) \subseteq M$ for some Hall π'-subgroup K of G.

Proof. Case 1: $\mathrm{Core}_G M = 1$. Since M is a π-solvable group, we have M contains Hall π'-subgroup of K. By Theorem 1.39 [1], we have

$$|G : K| = |G : M||M : K|.$$

Since $|G : M|$ is a π-number, we have K is a Hall π'-subgroup of G. Now we have $O_{\pi'}(G) \leq K \leq M$, so $O_{\pi'}(G) \leq \mathrm{Core}_G M = 1$. Since G is a π-solvable group, we have $O_\pi(G) \neq 1$. But G is a primitive group with a maximal subgroup M such that $\mathrm{Core}_G M = 1$. Therefore, for some $p \in \pi(G)$

$$N = O_p(G) = F(G) = C_G(O_p(G)) \neq 1, \ G = M[O_p(G)], \ \Phi(G) = 1$$

and $O_p(M) = 1$.

1. Let $F(M) \neq 1$, $q \in \pi(F(M))$, and let Q be a Sylow q-subgroup of M. Then $O_q(M) \neq 1$, $O_q(M) \subseteq Q$, and since $\mathrm{Core}_G M = 1$, we have $N_G(O_q(M)) = M$. Since $O_p(M) = 1$, it follows that p does not belong to $\pi(F(M))$ and $p \neq q$. The subgroup $D = N_G(Q) \cap O_p(G)$ is a normal subgroup of $N_G(Q)$ and

$$D \subseteq C_G(Q) \subseteq C_G(O_q(M)) \subseteq N_G(O_q(M)) = M.$$

Now $D = N_G(Q) \cap O_p(G) \subseteq M \cap O_p(G) = 1$. We consider the subgroup $L = N_G(Q)O_p(G)$. Since $N_G(Q) \cap O_p(G) = 1$, we have $L = [O_p(G)]N_G(Q)$. It follows from $G = [O_p(G)]M$ and Dedekind identity that

$$L = [O_p(G)]N_G(Q) = [O_p(G)](L \cap M),\ L/O_p(G) \simeq N_G(Q) \simeq L \cap M,$$

and $L \cap M$ is a q-closed subgroup. But $Q \subseteq L \cap M$, so Q is a normal subgroup of $L \cap M$. It follows from $|N_G(Q)| = |L \cap M|$ that $L \cap M = N_G(Q)$ and $N_G(Q) \subseteq M$.

2. Let $F(M) = 1$, and let K be a Hall π'-subgroup of M. By Lemma 2, $O_\pi(M)$ is a normal subgroup of G. Hence $O_\pi(M) \leq \mathrm{Core}_G M = 1$. Since G is a π-solvable group, we have $O_{\pi'}(M) \neq 1$ and $O_{\pi'}(M) \subseteq K$. Since $\mathrm{Core}_G M = 1$, we have $N_G(O_{\pi'}(M)) = M$. The subgroup $B = N_G(K) \cap O_p(G)$ is a normal subgroup of $N_G(K)$ and

$$B \subseteq C_G(K) \subseteq C_G(O_{\pi'}(M)) \subseteq N_G(O_{\pi'}(M)) = M.$$

Now $B = N_G(K) \cap O_p(G) \subseteq M \cap O_p(G) = 1$. We consider the subgroup $T = N_G(K)O_p(G)$. Since $N_G(K) \cap O_p(G) = 1$, we have $T = [O_p(G)]N_G(K)$. It follows from $G = [O_p(G)]M$ and Dedekind identity that

$$T = [O_p(G)]N_G(K) = [O_p(G)](T \cap M),\ T/O_p(G) \simeq N_G(K) \simeq T \cap M$$

and $T \cap M$ is a π'-closed subgroup. But Hall π'-subgroup K is contained in $T \cap M$, so K is a normal subgroup of $T \cap M$ and $T \cap M \subseteq N_G(K)$. It follows from isomorphism $N_G(K) \simeq T \cap M$ that $|N_G(K)| = |T \cap M|$, so $T \cap M = N_G(K)$ and $N_G(K) \subseteq M$. The Case 1 is proved.

Case 2: $N = \mathrm{Core}_G M \neq 1$. We consider the quotient group $\overline{G} = G/N$. Clearly, $\overline{G}$ is a π-solvable group and $|\overline{G} : \overline{M}| = |G : M|$ is a π-number, where $\overline{M} = M/N$. Since $\mathrm{Core}_{\overline{G}}\overline{M} = 1$, for the group $\overline{G}$ with a non-normal maximal subgroup $\overline{M}$, we apply Case 1.

1. Let $F(\overline{M}) \neq 1$, and let $q \in \pi(F(\overline{M}))$. By Case 1, $\overline{G}$ has a Sylow q-subgroup $\overline{Q}$ such that $N_{\overline{G}}(\overline{Q}) \subseteq \overline{M}$. Let $\overline{Q} = A/N$, and let Q be a Sylow q-subgroup of A. Then $QN/N = A/N = \overline{Q}$ and by Theorem 1.65 [1]

$$N_{\overline{G}}(\overline{Q}) = N_{G/N}(QN/N) = N_G(Q)N/N.$$

By the condition $N_G(Q)N/N \subseteq \overline{M} = M/N$, so $N_G(Q) \subseteq M$.

2. Let $F(M/\mathrm{Core}_G M) = 1$. By Case 1, $N_{\overline{G}}(\overline{K}) \subseteq \overline{M}$ for some Hall π'-subgroup $\overline{K}$ of $\overline{G}$. Let $\overline{K} = B/N$, and let R be a Hall π'-subgroup of B. It exists because B is a π-solvable group. Then R is a Hall π'-subgroup of G, $RN/N = B/N = \overline{K}$ and by Lemma 1,

$$N_G(R)N/N \subseteq N_{G/N}(RN/N) = N_{\overline{G}}(\overline{K}).$$

By induction, $N_G(R)N/N \subseteq \overline{M} = M/N$, so $N_G(R) \subseteq M$. The Theorem 3 is proved.

Note that Theorem 2 follows from Theorem 3 if $\pi = \{p\}$. If G is a solvable group, then in the condition of Theorem 3, $\pi(F(M/\mathrm{Core}_G M)) \neq \emptyset$, so Theorem 1 follows from Theorem 3.

PROOF OF COROLLARY 1.1. By Theorem 1, G has a Sylow q-subgroup Q such that $N_G(Q) \subseteq H$. Let H_1 be any subgroup such that $Q \subseteq H_1 \subseteq H$, and let $T = N_G(H_1)$. By the Frattini lemma, we have

$$T = N_T(Q)H_1 \subseteq N_G(Q)H_1 \subseteq H, \; T = N_G(H_1) \subseteq H.$$

The Corollary is proved.

References

[1] Monakhov V. S. Introduction to the Theory of Finite Groups and Their Classes. Minsk: High school. 2006. (In Russian)

[2] Vedernikov V. A. About π-properties of a finite groups // In the book: Arithmetic and subgroups a structure of finite groups. Minsk: Science and technics. 1986. P. 13–19. (In Russian)

D. V. GRITSUK

Department of mathematics, Gomel F. Scorina State University, Gomel 246019, BELARUS

E-mail address: Dmitry.Gritsuk@gmail.com

V. S. MONAKHOV

Department of mathematics, Gomel F. Scorina State University, Gomel 246019, BELARUS

E-mail address: Victor.Monakhov@gmail.com